

Integral Representation and Embedding of Weak Markov Systems

R. A. ZALIK

*Division of Mathematics, Department of Algebra,
Combinatorics and Analysis, Auburn University, Auburn, Alabama 36849-5307, U.S.A.*

Communicated by Oved Shisha

Received September 16, 1982; revised February 12, 1988

In the sequel, A will always denote a subset of the real line having at least $n+2$ elements ($n \geq 0$), $l_1 = \inf(A)$, $l_2 = \sup(A)$, and $I(A)$ will denote the convex hull of A (thus for example, if $A = [2, 3) \cup (4, \infty)$, then $I(A) = [2, \infty)$); $Z_n = \{z_0, \dots, z_n\}$ will be a set of real valued functions defined on A ; by $S(Z_n)$ we shall denote the linear span of Z_n . We shall call Z_n a weak Čebyšev system (Čebyšev system), provided that Z_n be linearly independent on A , and for every choice of $n+1$ points t_i of A , with $t_0 < t_1 < \dots < t_n$, $\det[z_i(t_j); i, j = 0, \dots, n] \geq 0$ (> 0). If Z_k is a (weak) Čebyšev system for $k = 0, \dots, n$, then Z_n will be called a (weak) Markov system. A normalized—or normed—(weak) Markov system is a (weak) Markov system Z_n for which $z_0 \equiv 1$. Markov systems are also called complete Čebyšev systems (cf. Karlin and Studden [2]). We shall say that $U_n = \{u_0, \dots, u_n\}$ has been obtained from Z_n by a triangular linear transformation if $u_0 = z_0$, and

$$u_k - z_k \in S(Z_{k-1}), \quad k = 1, 2, \dots, n.$$

Note that if Z_n is linearly independent then, for $k = 0, 1, \dots, n$, U_k is a basis of $S(Z_k)$. We shall adopt the convention that if $b \leq a$, then $[a, b) = \emptyset$.

In [6, Theorem 1] we gave an integral representation of Markov systems. Recently Zielke [11] gave a counterexample and a corrected version of this result, and generalized it to a class of normalized weak Markov systems. The purpose of our paper is to extend the results of [11], using a refinement of a new embedding property of normalized weak Markov systems developed in [7].

A system Z_n will be called nondegenerate if for every c in A , Z_n is linearly independent both on $(-\infty, c) \cap A$ and on $(c, \infty) \cap A$, and it will be called weakly nondegenerate provided that the following conditions are satisfied:

Condition I. For every real number c , Z_n is linearly independent on at least one of the sets $(c, \infty) \cap A$ and $(-\infty, c) \cap A$.

Condition E. For every point c in $I(A)$ we have:

(a) If Z_n is linearly independent on $[c, \infty) \cap A$, then there exists a set U_n , obtained from Z_n by a triangular linear transformation, such that for any sequence $\{k(r); r=0, \dots, m\}$ with $k(0) \geq 0$ and $k(m) \leq n$ that is either strictly increasing or contains exactly one element, the set $\{u_{k(r)}; r=0, \dots, m\}$ is a weak Markov system on $[c, \infty) \cap A$.

(b) If Z_n is linearly independent on $(-\infty, c] \cap A$, then there exists a set V_n , obtained from Z_n by a triangular linear transformation, such that for every sequence $\{k(r); r=0, \dots, m\}$ with $k(0) \geq 0$ and $k(m) \leq n$ that is either strictly increasing or contains exactly one element, $\{(-1)^{r-k(r)}v_{k(r)}; r=0, \dots, m\}$ is a weak Markov system on $(-\infty, c] \cap A$.

Finally Z_n will be called "representable" if for any point c in A there exist a set $U_n = \{u_0, \dots, u_n\}$, obtained from Z_n by a triangular linear transformation; a strictly increasing and bounded real function $h(t)$, defined on A and such that $h(c) = c$; and continuous, increasing, and nonconstant real functions $w_k(t)$, defined on $I(h(A))$, such that for all x in A

$$\begin{aligned} u_0 &\equiv 1 \\ u_1(x) &= \int_c^{h(x)} dw_1(t_1) \\ &\vdots \\ u_n(x) &= \int_c^{h(x)} \int_c^{t_1} \cdots \int_c^{t_{n-1}} dw_n(t_n) \cdots dw_1(t_1). \end{aligned} \tag{1}$$

In [11], Zielke essentially proved that a nondegenerate normalized weak Markov system is representable. Our main result is:

THEOREM 1. *Every weakly nondegenerate normalized weak Markov system is representable.*

Remarks. (i) In the statement of [11, Theorem 3], Zielke asserts that the representation (1) is valid for *some* point c , but in the proof of the theorem he actually shows that a representation exists for *any* point c in A . The distinction is, however, immaterial: If (1) is satisfied for some point c in A it is easy to see that for any other point c' in A there is a basis U'_n of $S(Z_n)$, obtained from U_n by a triangular linear transformation, having a representation of the form (1) with c replaced by c' .

(ii) It can be shown that every nondegenerate normalized weak Markov system satisfies Condition E (this has essentially been done in the proof of

[8, Theorem 2]). Thus, every nondegenerate normalized weak Markov system is weakly nondegenerate. The converse, however, is false: Let $u_0 \equiv 1$, $u_1(x) = x$ on $(0, 1]$, $u_1(x) = 1$ on $[1, 2)$, and $u_2(x) = [u_1(x)]^2$ on $(0, 2)$. Then $U_2 = \{u_0, u_1, u_2\}$ is a weakly nondegenerate normalized weak Markov system on $(0, 2)$. However U_2 is not nondegenerate there.

(iii) The converse of Theorem 1 is false: let $h(t) = t$, $w_1(t) = 1$ on $(-1, 0)$, $w_1(t) = t + 1$ on $[0, 1)$, $w_2(t) = t$ on $(-1, 0)$, and $w_2(t) = 0$ on $[0, 2)$. If $U_2 = \{u_0, u_1, u_2\}$ has a representation of the form (1), it is readily seen that $u_2 \equiv 0$ on $(-1, 1)$.

We shall call Z_n strongly representable if it is representable and all the functions $w_i(t)$ are strictly increasing on $h(A)$. We shall say that A has property (B), if for any two elements of A there is a third element of A between them. As a consequence of Theorem 1 we shall prove:

THEOREM 2. *Let A have property (B) and assume that Z_n is weakly nondegenerate. Then Z_n is a normalized Markov system if and only if it is strongly representable.*

COROLLARY. *Let A have property (B) and assume that Z_n is weakly nondegenerate. Then if Z_n is a normalized Markov system on A there is a function z_{n+1} such that also $Z_n \cup \{z_{n+1}\}$ is a normalized Markov system on A .*

If Z_n is a Markov system, it is obvious that Condition I is satisfied. If, moreover, A satisfies property (B), it is easy to see that Condition E is satisfied for any point in $(l_1, l_2) \cap A$, making the assumption of weak nondegeneracy redundant if neither l_1 nor l_2 are in A ; thus this corollary generalizes the main result of [9]. Since it is not known as yet under what circumstances Condition E will be satisfied at an endpoint, it is at present unclear whether the corollary also generalizes the main result of [5]. We intend to study this problem in a later paper.

A system Z_n is called C -bounded if every element of Z_n is bounded on the intersection of A with any compact subset of $I(A)$; if A is an interval and every element of Z_n is absolutely continuous in any closed subinterval of A , we shall say that Z_n is C -absolutely continuous. If $V_n = \{v_0, \dots, v_n\}$ is a set of real functions defined on a real set B we say that Z_n can be embedded in V_n if there is a strictly increasing function $h: A \rightarrow B$ such that $v_i[h(t)] = z_i(t)$ for every $t \in A$ and $i = 0, 1, \dots, n$. The function h is called an embedding function.

In the proof of Theorem 1 we shall need the following refinement of the theorem of [7]:

THEOREM 3. *Let c be an element of A . If Z_n is a weakly nondegenerate normalized weak Markov system on A , then Z_n can be embedded in a weakly nondegenerate normalized weak Markov system V_n of C -absolutely con-*

tinuous functions defined on an open bounded interval, and V_n and the embedding function $h(t)$ can be chosen so that $h(c)=c$. Moreover if A satisfies property (B), the converse statement is also true.

The proof of Theorem 3 is based on the following auxiliary propositions:

LEMMA 1. Let Z_n be a weakly nondegenerate weak Markov system on a set A , let $p: A \rightarrow R$ be a strictly increasing function, and let $v_r(t) = z_r(p^{-1}(t))$, $r=0, \dots, n$. Then V_n is a weakly nondegenerate weak Markov system on $p(A)$.

The proof of Lemma 1 is straightforward and will be omitted.

LEMMA 2. Let $[a, b]$ be a closed bounded interval. Assume that f is a continuous function of bounded variation and that g is a strictly increasing continuous function, both defined on $[a, b]$. For $a \leq \alpha \leq \beta \leq b$, let $V(f, \alpha, \beta)$ denote the total variation of f on $[\alpha, \beta]$. Let $c \in [a, b]$ be arbitrary but fixed, and define $v(f, t)$ to equal $V(f, c, t)$ on $[c, b]$ and $-V(f, t, c)$ on $[a, c]$. Finally, let $q(t) = g(t) + v(f, t)$ and $h(t) = f[q^{-1}(t)]$. Then $h(t)$ is absolutely continuous on $[q(a), q(b)]$.

Proof of Lemma 2. The hypotheses imply that $q(t)$ is strictly increasing and continuous; thus $q^{-1}(t)$ is strictly increasing on $[q(a), q(b)]$. If $a < s_1 < s_2 < b$, then $|f(s_2) - f(s_1)| \leq V(f, s_1, s_2) = v(f, s_2) - v(f, s_1) \leq v(f, s_2) - v(f, s_1) + g(s_2) - g(s_1) = q(s_2) - q(s_1)$. Thus, if $(\alpha_1, \beta_1), (\alpha_2, \beta_2), \dots, (\alpha_n, \beta_n)$ are disjoint subintervals of $[q(a), q(b)]$ we have

$$\begin{aligned} \sum_{i=1}^n |h(\beta_i) - h(\alpha_i)| &= \sum_{i=1}^n |f[q^{-1}(\beta_i)] - f[q^{-1}(\alpha_i)]| \\ &\leq \sum_{i=1}^n (q[q^{-1}(\beta_i)] - q[q^{-1}(\alpha_i)]) = \sum_{i=1}^n (\beta_i - \alpha_i), \end{aligned}$$

and the conclusion follows.

Q.E.D.

The following lemma implies that every weakly nondegenerate normalized weak Markov system is C -bounded:

LEMMA 3. Let $U_n = \{u_0, \dots, u_n\}$ be a weakly nondegenerate normalized weak Markov system on a set A , let $I_1 = \inf(A)$, $I_2 = \sup(A)$, $c \in I(A)$, and let u be any function in $S(U_n)$.

(a) If $c > I_1$ and c is a point of accumulation of $(I_1, c) \cap A$, then $\lim_{t \rightarrow c^-} u(t)$ exists and is finite.

(b) If $c < I_2$ and c is a point of accumulation of $(c, I_2) \cap A$, then $\lim_{t \rightarrow c^+} u(t)$ exists and is finite.

Proof. We only prove (a); the proof of (b) is similar and will be omitted.

We proceed by induction. The assertion is trivially true for $n=0$. To prove the inductive step, assume that for any function w in $S(U_{n-1})$ (where $U_{n-1} = \{u_0, \dots, u_{n-1}\}$), $\lim_{t \rightarrow c^-} w(t)$ exists and is finite. If U_n is linearly independent on $(-\infty, c) \cap A$ there is a number $d \in A$ such that $d \geq c$. Indeed, this is obvious if $c < l_2$, whereas if $c = l_2$ we can take $d = l_2$. Since clearly U_n is linearly independent on $(-\infty, d] \cap A$, by Condition E we conclude that there is a function $u = u_n + w$, with $w \in S(U_{n-1})$, such that u is monotonic on $(-\infty, d] \cap A$, whence the conclusion readily follows. Assume now that U_n is linearly dependent on $(-\infty, c) \cap A$. Condition I then implies that U_n is linearly independent on $(c, \infty) \cap A$, and therefore on any set of the form $(d', \infty) \cap A$, $d' < c$. Another application of Condition E readily yields the conclusion for this case as well. Q.E.D.

The proof of the next proposition was sketched in [7].

LEMMA 4. *Let Z_n be a normalized weak Markov system of bounded functions defined on a closed interval $I = [a, b]$. Then all the elements of $S(Z_n)$ are of bounded variation on I .*

Proof. Let z be a function in $S(Z_n)$, arbitrary but fixed, let γ be any real number, and let $v(\gamma)$ denote the number of sign changes of $z(t) - \gamma$. Since [10, p. 12, Lemma 4.1] implies that $v(\gamma) \leq n$, and the boundedness of $z(t)$ implies that $v(\gamma)$ has bounded support, the conclusion follows from, e.g., [4, p. 257, Theorem 6]. Q.E.D.

LEMMA 5. *Let Z_n be a weakly nondegenerate normalized weak Markov system defined on an interval I (open, closed, or semiopen), and let $c \in I$. If z_1 is continuous at c , then all the elements of $S(Z_n)$ are continuous at c .*

Proof. We shall only prove that if $c > \inf(I)$, then all the elements of $S(Z_n)$ are left-continuous at c . The proof of the other case is similar and will be omitted.

We proceed by induction on n . For $n=1$ the assertion is true by hypothesis; assume therefore that $n > 1$.

If Z_n is linearly independent on $S_1 = (-\infty, c] \cap I$, then from Condition E we readily conclude that there is a set U_n , obtained from Z_n by a triangular linear transformation, such that both $\{1, (-1)^{n-1}u_n\}$ and $\{1, u_1, (-1)^nu_n\}$ are weak Markov systems on S_1 . The first assertion is equivalent to saying that $(-1)^{n-1}u_n$ is increasing on S_1 , from which we conclude that $(-1)^{n-1}u_n(c^-) \leq (-1)^{n-1}u_n(c)$. The linear independence implies that there is a point t_0 in $(-\infty, c) \cap I$ such that $u_1(t_0) < u_1(c)$.

Assume that $t_0 < t < c$; then, subtracting the second column from the third, we have

$$\begin{aligned} 0 &\leq \begin{vmatrix} 1 & 1 & 1 \\ u_1(t_0) & u_1(t) & u_1(c) \\ (-1)^n u_n(t_0) & (-1)^n u_n(t) & (-1)^n u_n(c) \end{vmatrix} \\ &= (-1)^n \begin{vmatrix} 1 & 1 & 0 \\ u_1(t_0) & u_1(t) & u_1(c) - u_1(t) \\ u_n(t_0) & u_n(t) & u_n(c) - u_n(t) \end{vmatrix}. \end{aligned}$$

Since $u_1(t)$ is continuous at c , passing to the limit we have

$$\begin{aligned} 0 &\leq (-1)^n \begin{vmatrix} 1 & 1 & 0 \\ u_1(t_0) & u_1(c) & 0 \\ u_n(t_0) & u_n(c^-) & u_n(c) - u_n(c^-) \end{vmatrix} \\ &= (-1)^n [u_1(c) - u_1(t_0)][u_n(c) - u_n(c^-)], \end{aligned}$$

whence we conclude that $(-1)^{n-1} u_n(c) \leq (-1)^{n-1} u_n(c^-)$. We have therefore shown that $u_n(t)$ is left-continuous at c . Since $u_n = z_n + w$, with $w \in S(Z_{n-1})$, applying the inductive hypothesis we conclude that also z_n is left-continuous at c .

If Z_n is linearly dependent on $(-\infty, c] \cap I$, from Condition I we conclude that Z_n must be linearly independent on $(c, \infty) \cap I$. Thus, if d is an arbitrary but fixed point in $(-\infty, c) \cap I$, it is clear that Z_n is linearly independent on $J_2 = [d, \infty) \cap I$, whence by Condition E there is a set V_n , obtained from Z_n by a triangular linear transformation, such that both $\{1, v_n\}$ and $\{1, v_1, v_n\}$ are normalized weak Markov systems on J_2 . The first assertion is equivalent to saying that v_n is increasing on J_2 , from which we conclude that $v_n(c^-) \leq v_n(c)$. The linear independence implies that there is a point t_1 in $(c, \infty) \cap I$ such that $v_1(c) < v_1(t_1)$. Choosing $t < c$ and proceeding as in the preceding paragraph, we deduce that $[v_n(c^-) - v_n(c)][v_1(t_1) - v_1(c)] \geq 0$, whence $v_n(c) \leq v_n(c^-)$, and the conclusion readily follows. Q.E.D.

Proof of Theorem 3. Assume that Z_n is a weakly nondegenerate normalized weak Markov system.

From [7] we know that Z_n can be embedded in a normalized weak Markov system $U_n = \{u_0, \dots, u_n\}$ of continuous functions defined on an open interval (a_1, b_1) , and such that if h is the embedding function then $h(c) = c$.

Assuming now that Z_n is weakly nondegenerate, we shall adapt the procedure outlined in the proof of the theorem of [7] to show that U_n is also weakly nondegenerate.

Let $S = \{s_i\}$ denote the set of points of accumulation of A at which $z_1(t)$ has jump discontinuities. If $s_i \in S \cap (l_1, l_2)$, let $d_i = 2^{-i}$; on the other hand if $s_i \in A$, let $a_i = 2^{-(i+1)}$ if $z_1(s_i^+) - z_1(s_i) \neq 0$, and 0 otherwise. If $s_i = l_1$ and $l_1 \in A$, let $d_i = a_i = |z_1(s_i^+) - z_1(s_i)|$, whereas if $s_i = l_2$ and $l_2 \in A$, we define $a_i = |z_1(s_i) - z_1(s_i^-)|$. Let $q(t) = t + \sum_{s_j < t} d_j$ if $t \in A$ but $t \notin S$, whereas for $t_i \in A \cap S$ we define $q(t_i) = t_i + (\sum_{s_j < t_i} d_j) + a_i$. It is clear that $q(t)$ is strictly increasing. (Note that there is a typographical error in the definitions of α_i and β_i in [7]. They should be defined in a manner similar to that of a_i above.)

Setting $z_k^{(0)}(t) = z_k[q^{-1}(t)]$, we infer from Lemma 1 that $Z_n^{(0)}$ is weakly nondegenerate on $A^{(0)} = q(A)$. Moreover, it has the property that $z_1^{(0)}$ is either continuous or has a removable discontinuity at every point of accumulation of $A^{(0)}$.

Let $l_1^{(0)} = \inf(A^{(0)})$, $l_2^{(0)} = \sup(A^{(0)})$. If $l_1^{(0)}$ belongs to $A^{(0)}$, define $z_r^{(1)}$ to equal $z_r^{(0)}(l_1)$ on $(-\infty, l_1^{(0)})$; if $l_2^{(0)}$ belongs to $A^{(0)}$, define $z_r^{(1)}$ to equal $z_r^{(0)}(l_2^{(0)})$ on $(l_2^{(0)}, \infty)$; moreover, let $z_r^{(1)} = z_r^{(0)}$ on $A^{(0)}$. Clearly $Z_n^{(1)}$ is a weakly nondegenerate normalized weak Markov system defined on a set $A^{(1)}$ that has no first nor last element. Let $l_1^{(1)} = \inf(A^{(1)})$, $l_2^{(1)} = \sup(A^{(1)})$, and let $\bar{A}^{(1)}$ denote the closure of $A^{(1)}$ in the relative topology of $I = (l_1^{(1)}, l_2^{(1)})$. If x is in $\bar{A}^{(1)}$ but not in $A^{(1)}$, define $z_r^{(2)}(x) = \lim_{t \rightarrow x^-} z_r^{(1)}(t)$, $r = 0, \dots, n$, if x is a point accumulation of $(-\infty, x) \cap A^{(1)}$, or $z_r^{(2)}(x) = \lim_{t \rightarrow x^+} z_r^{(1)}(t)$, $r = 0, \dots, n$, if it is not (this can be done because of Lemma 3), whereas for x in $A^{(1)}$, let $z_r^{(2)}(x) = z_r^{(1)}(x)$. Clearly $Z_n^{(2)}$ is a normalized weak Markov system on $\bar{A}^{(1)}$.

It is readily seen that $Z_n^{(2)}$ is weakly nondegenerate on $\bar{A}^{(1)}$. To prove Condition I for $Z_n^{(2)}$, assume, e.g., that $Z_n^{(2)}$ is linearly dependent on $[c, \infty) \cap \bar{A}^{(1)}$. From Condition I for $Z_n^{(1)}$ we readily infer that $Z_n^{(1)}$ is linearly independent on $(-\infty, c] \cap A^{(1)}$, which clearly implies that $Z_n^{(2)}$ is linearly independent on $(-\infty, c] \cap \bar{A}^{(1)}$. To prove Condition E, assume, for example, that $Z_n^{(2)}$ is linearly independent on $[c, \infty) \cap \bar{A}^{(1)}$. Let $d \in A^{(1)}$, $d < c$ be arbitrary but fixed. Since $Z_n^{(1)}$ is clearly linearly independent on $[d, \infty) \cap A^{(1)}$, applying Condition E to $Z_n^{(1)}$ on $[d, \infty) \cap A^{(1)}$ and passing to the limit, Condition E for $Z_n^{(2)}$ on $[c, \infty) \cap \bar{A}^{(1)}$ readily follows.

Clearly the complementary set of $\bar{A}^{(1)}$ in $(l_1^{(1)}, l_2^{(1)})$, if not empty, is a disjoint union of open intervals V_j ; moreover if $c_j = \inf(V_j)$ and $d_j = \sup(V_j)$, then both c_j and d_j are in $\bar{A}^{(1)}$. Let w_r be defined on I as follows: If $t \in \bar{A}^{(1)}$, then $w_r(t) = z_r^{(2)}(t)$. On the other hand, if $t \notin \bar{A}^{(1)}$, then $c_i < t < d_i$ for some i . In this case, define $w_r(t) = [(d_i - t) z_r^{(2)}(c_i) + (t - c_i) z_r^{(2)}(d_i)] / (d_i - c_i)$. (Note that $t = [(d_i - t)c_i + (t - c_i)d_i] / (d_i - c_i)$.) It is readily seen that $W_n = \{w_0, \dots, w_n\}$ is a normalized weakly nondegenerate weak Markov system, defined on the open interval I . Since $z_1^{(2)}$ is clearly continuous on $\bar{A}^{(1)}$, and w_1 is obtained from it by linear interpolation, we readily deduce that w_1 is continuous on I . Applying Lemma 5, we thus conclude that all

the elements of W_n are continuous on I . We have therefore shown that Z_n can be embedded in a weakly nondegenerate normalized weak Markov system W_n of continuous functions defined on an open interval I . Moreover, from Lemma 4 we know that the elements of $S(W_n)$ are of bounded variation on every closed subinterval of I . Thus, if the functions $v(w_k, t)$ are defined as in Lemma 2, $p(t) = t + \sum_{k=1}^n v(w_k, t)$, $I_1 = p(I)$, $w_i^{(1)}(t) = w_i[p^{-1}(t)]$, $i = 0, 1, 2, \dots, n$, and $W_n^{(1)} = \{w_0^{(1)}, w_1^{(1)}, \dots, w_n^{(1)}\}$, it is readily seen from Lemmas 1 and 2 that $W_n^{(1)}$ is a weakly nondegenerate normalized weak Markov system of C -absolutely continuous functions on I_1 . Thus, there is a strictly increasing function $h(t)$ that embeds Z_n into $W_n^{(1)}$. Setting $p_1(t) = c - h(c) + t$, $h_1(t) = p_1[h(t)]$, $w_i^{(2)}(t) = w_i^{(1)}[p_1^{-1}(t)]$, and $W_n^{(2)} = \{w_0^{(2)}, \dots, w_n^{(2)}\}$, it is easy to see that $h_1(t)$ embeds Z_n into $W_n^{(2)}$, and that $h_1(c) = c$. Making if necessary a change of variable of the form $c + \arctan(t - c)$ to ensure the boundedness of the domain of the elements of $W_n^{(2)}$, the conclusion readily follows.

The proof of the converse is trivial and will be omitted.

Q.E.D.

To prove Theorem 1 we also need the following:

LEMMA 6. *Let $U_n = \{u_0, \dots, u_n\}$ be a weakly nondegenerate weak Markov system on an interval (a, b) . If for some c in (a, b) , $u_0(c) = 0$, then $u_k(c) = 0$, $k = 1, 2, \dots, n$.*

Proof. We proceed by induction on n . For $n = 0$ the assertion is true by hypothesis. To prove the inductive step, assume first that U_n is linearly independent on (a, c) . Then from Condition E we readily conclude that there is a set V_n , obtained from U_n by a triangular linear transformation, such that both $\{(-1)^n v_n\}$ and $\{v_0, (-1)^{n+1} v_n\}$ are weak Markov systems on $(a, c]$. From the first condition we conclude that $(-1)^n v_n(c) \geq 0$. The linear independence implies that there is a point $t_0 \in (a, c)$, such that $v_0(t_0) \neq 0$. Applying the second condition we readily deduce that $v_0(t_0) > 0$ and that $v_0(t_0)(-1)^{n+1} v_n(c) \geq 0$. Thus $(-1)^n v_n(c) \leq 0$, and the assertion readily follows.

If U_n is linearly dependent on (a, c) , from Condition I we deduce that it must be linearly independent on (c, b) , and the assertion is proved by a similar procedure.

Q.E.D.

Proof of Theorem 1. Let Z_n be a weakly nondegenerate normalized weak Markov system. Without loss of generality we can assume that $z_i(c) = 0$, $i = 1, \dots, n$. From Theorem 3 we know that there is a strictly increasing function $p: A \rightarrow (a_1, b_1)$ and a weakly nondegenerate normalized weak Markov system $\{q_0, \dots, q_n\}$ of C -absolutely continuous functions defined on (a_1, b_1) , such that $z_i = q_i \circ p$, $i = 0, \dots, n$, and $p(c) = c$. Clearly $q_i(c) = 0$, $i = 1, \dots, n$; moreover, if D is the set of points on which the func-

tions q_i are differentiable, then the measure of D equals $b_1 - a_1$. Let I be a subinterval of (a_1, b_1) , and let $\{k(r); r=0, \dots, m\}$ be a strictly increasing sequence with $k(0)=0$, $k(1) \geq 1$, and $k(m) \leq n$. Since the functions $q_i(t)$ are C -absolutely continuous, it is readily seen that $\{q'_{k(r)}; r=1, \dots, m\}$ is linearly dependent on $I \cap D$ if and only if $\{q_{k(r)}; r=0, \dots, m\}$ is linearly dependent on I . Thus, proceeding as in [10, Theorem 11.3(b)] we readily infer that $Q'_{n-1} = \{q'_1, \dots, q'_n\}$ is a weakly nondegenerate weak Markov system on D ; thus if S is the subset of D on which $q'_i \neq 0$, and $m_i = q'_i/q'_1$, from Lemma 6 we readily deduce that $M_{n-1} = \{m_1, \dots, m_n\}$ is a weakly nondegenerate normalized weak Markov system on S .

Let $a_2 = \inf(S)$, $b_2 = \sup(S)$, and assume for instance that $a_2 > a_1$ and $b_2 < b_1$. This implies that $q'_i(t) = 0$ on $(a_1, a_2) \cap D$ and on $(b_2, b_1) \cap D$. In particular, Condition I implies that Q'_{n-1} is linearly independent on $(a_1, b_2] \cap D$. Thus Condition E implies that there is a system $R_{n-1} = \{r_1, \dots, r_n\}$, obtained from Q'_{n-1} by a triangular linear transformation, such that $\{(-1)^{i-1}r_i\}$ and $\{r_i, (-1)^i r_i\}$, $i=2, \dots, n$, are weak Markov systems on D . Since $r_1 \equiv q'_1 > 0$ on S , this means that $(-1)^i r_i/r_1$ is both increasing and nonpositive (and therefore bounded from above) on S . Setting $u_i = r_i/r_1$ on S and $u_i(t) = \lim_{t \rightarrow b_2^-} r_i(t)/r_1(t)$ on $[b_2, b_1)$, it is clear that $U_n = \{u_1, \dots, u_n\}$ is a weakly nondegenerate weak Markov system on $S \cup [b_2, b_1)$. This means that there is a weakly nondegenerate normalized weak Markov system $M_{n-1}^{(0)} = \{m_1^{(0)}, \dots, m_n^{(0)}\}$ on $S \cup [b_2, b_1)$ that coincides with M_{n-1} on S , and such that the functions $m_k^{(0)}$ are constant on $[b_2, b_1)$. Applying Condition E again and using a similar procedure, it is easy to see that $M_{n-1}^{(0)}$ can be extended to the left; i.e., there exists a weakly nondegenerate normalized weak Markov system $M_{n-1}^{(1)} = \{m_1^{(1)}, \dots, m_n^{(1)}\}$ that coincides with M_{n-1} on S , and such that the functions $m_k^{(1)}$ are constant on $(a_1, a_2]$ and on $[b_2, b_1)$. Proceeding as in the proof of Theorem 3 we readily see that there is a weakly nondegenerate normalized weak Markov system $V_{n-1} = \{v_1, \dots, v_n\}$ on (a_1, b_1) that coincides with M_{n-1} on S . Since Lemma 6 implies that all the functions q'_i vanish on $D - S$, we conclude that $q'_i(t) = q'_1(t) v_i(t)$ for every t in D and $i=1, \dots, n$. It is therefore clear that for every x in (a_1, b_1) ,

$$q_i(x) = \int_c^x q'_1(t) v_i(t) dt, \quad i=1, \dots, n. \quad (2)$$

The proof is completed by induction. For $n=1$ the assertion of the theorem follows from (2). We now proceed to the proof of the inductive step.

By inductive hypothesis there is a basis $\{\bar{v}_1, \dots, \bar{v}_n\}$ of the linear span of $\{v_1, \dots, v_n\}$, such that for $i=1, \dots, n$ and $x \in (a_1, b_1)$, $\bar{v}_i(x) = p_i[h(x)]$, where

$$p_i(x) = \int_c^x \int_c^{t_2} \dots \int_c^{t_{i-1}} dw_i(t_i) \dots dw_2(t_2),$$

$h(x)$ is strictly increasing and bounded on (a_1, b_1) , $h(c) = c$, and the functions w_i are continuous and increasing on $(h(a_1^+), h(b_1^-))$. There is no loss of generality if we assume that $\bar{v}_i = v_i$, $i = 1, \dots, n$. It is clear that the inverse function of h can be extended to an increasing (but not necessarily strictly increasing) function g , continuous on $(h(a_1^+), h(b_1^-))$; thus since the functions $p_i(x)$ are continuous, setting $w_1(x) = \int_c^{g(x)} q_1'(t) dt$ and applying [1, p. 182, Lemma 8(f); 3, p. 368, Theorem 1] we easily conclude that $q_i(x) = \int_c^{h(x)} p_i(t) dw_1(t)$, whence the assertion readily follows. Q.E.D.

Proof of Theorem 2. From Theorem 1 we know that Z_n is representable. Let $U_n = \{u_0, \dots, u_n\}$ be a basis of $S(Z_n)$ having the representation (1). One easily sees (as in [5, Lemma 2]) that U_n is a normalized Markov system on A .

Assume that for some k , $w_k(t)$ is constant on some subinterval I of $I(h(A))$ that contains two points of $h(A)$. By an inductive procedure involving the number of integrations we see that $u_k[h^{-1}(t)]$ can be expressed as a linear combination of $u_0(h^{-1}(t)), \dots, u_{k-1}(h^{-1}(t))$ on I . Thus u_k can be expressed as a linear combination of u_0, \dots, u_{k-1} on $h^{-1}(I)$. Since h is strictly increasing and A has property (B), $h^{-1}(I)$ has an infinite number of points. Since U_k is a Čebyšev system we have obtained a contradiction.

To prove the converse, let U_n be a basis having a representation of the form (1), where the functions h and w_i satisfy the hypotheses of the theorem. For $k = 0, \dots, n$, let $v_k(x) = u_k[h^{-1}(x)]$; it suffices to prove that V_n is a normalized Markov system on $h(A)$. Since h is strictly increasing, it is clear $h(A)$ has property (B). Thus, if $\{x_i; i = 0, \dots, n\}$ is an arbitrary subset of $h(A)$, with $x_0 < x_1 < \dots < x_n$, there is a subset $\{t_i; i = 1, \dots, n\}$ of $h(A)$ with $x_{i-1} < t_i < x_i$. We now proceed by induction. The assertion is clearly trivial for $n = 0$ and $n = 1$. To prove the inductive step, let $a = \inf(h(A))$, $b = \sup(h(A))$, and let $f_r(t)$ be defined as follows: $f_1 = 1$, $f_2(t) = \int_c^t dw_2(t_1)$, and for $r = 3, \dots, n$, $f_r(t) = \int_c^t \int_c^{t_2} \dots \int_c^{t_{r-1}} dw_r(t_r) \dots dw_2(t_2)$. Clearly $v_r(x) = \int_c^x f_r(t) dw_1(t)$, $r = 1, \dots, n$. By inductive hypothesis $\{f_1, \dots, f_n\}$ is a normalized Markov system on $h(A)$. In particular, this implies that for every k , $k = 1, \dots, n$, $\det[f_i(t_j); i, j = 1, \dots, k] > 0$. By continuity we conclude that for each i there is a subinterval J_i of (x_{i-1}, x_i) such that if $s_i \in J_i$ for each i , then for each k , $k = 1, \dots, n$, $\det[f_i(s_j); i, j = 1, \dots, k] > 0$. Proceeding as in, e.g., the proof of [2, p. 382, Lemma 1], we see that for any k , $k = 0, 1, \dots, n$,

$$\det[v_i(x_j); i, j = 0, \dots, k] \\ = \int_{x_0}^{x_1} \int_{x_1}^{x_2} \dots \int_{x_{k-1}}^{x_k} \det[f_i(s_j); i, j = 1, \dots, k] dw_1(s_k) dw_1(s_{k-1}) \dots dw_1(s_1).$$

Since, moreover, $\{f_1, \dots, f_n\}$ is a weak Markov system on (a, b) , the conclusion readily follows. Q.E.D.

Proof of corollary. From Theorem 2 there is a basis U_n of Z_n having a representation of the form (1), where $h(t)$ is strictly increasing, and the $w_i(t)$ are increasing on $I(h(A))$, and strictly increasing on $h(A)$. Setting $w_{n+1}(x) = x$ and

$$z_{n+1}(x) = \int_c^{h(x)} \int_c^{t_1} \cdots \int_c^{t_n} dw_{n+1}(t_{n+1}) \cdots dw_1(t_1),$$

we readily obtain the conclusion.

Q.E.D.

ACKNOWLEDGMENTS

The author is grateful to C. E. Linderholm, R. N. Mera Valverde, R. Zielke, and D. Zwick for their helpful comments, and in particular to T. A. Kilgore for his critical reading of several versions of the manuscript.

REFERENCES

1. N. DUNFORD AND J. T. SCHWARTZ, "Linear Operators," Part I, Interscience, New York, 1957.
2. S. KARLIN AND W. J. STUDDEN, "Tchebycheff Systems: With Applications in Analysis and Statistics," Interscience, New York, 1966.
3. A. N. KOLMOGOROV AND S. V. FOMIN, "Introductory Real Analysis," Dover, New York, 1975.
4. I. J. SCHOENBERG, On variation diminishing approximation methods, in "On Numerical Approximation" (R. E. Langer, Ed.), pp. 249–274, Univ. of Wisconsin Press, Madison, 1959.
5. R. A. ZALIK, Existence of Tchebycheff extensions, *J. Math. Anal. Appl.* **51** (1975), 68–75.
6. R. A. ZALIK, Integral representation of Tchebycheff systems, *Pacific J. Math.* **68** (1977), 553–568.
7. R. A. ZALIK, Embedding of weak Markov systems, *J. Approx. Theory* **41** (1984), 253–256; Erratum, *J. Approx. Theory* **43** (1985), 396.
8. R. A. ZALIK AND D. ZWICK, On extending the domain of definition of Čebyšev and weak Čebyšev systems, *J. Approx. Theory* **57** (1989), 202–210.
9. R. ZIELKE, Alternation properties of Tchebycheff-systems and the existence of adjointed functions, *J. Approx. Theory* **10** (1974), 172–184.
10. R. ZIELKE, Discontinuous Čebyšev systems, in "Lecture Notes in Mathematics," Vol. 707, Springer-Verlag, Berlin/New York, 1979.
11. R. ZIELKE, Relative differentiability and integral representation of a class of weak Markov systems, *J. Approx. Theory* **44** (1985), 30–42.